



# UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office  
Address: COMMISSIONER OF PATENTS AND TRADEMARKS  
Washington, D.C. 20231  
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
08/785,532	01/17/1997	JOE W. GRAY	2500.124US2	4124

22798 7590 03/11/2003

QUINE INTELLECTUAL PROPERTY LAW GROUP, P.C.  
P O BOX 458  
ALAMEDA, CA 94501

EXAMINER

DAVIS, MINH TAM B

ART UNIT PAPER NUMBER

1642

DATE MAILED: 03/11/2003

35

Please find below and/or attached an Office communication concerning this application or proceeding.

**Office Action Summary**

Application No.

08/785,532

Applicant(s)

GRAY ET AL.

Examiner

MINH-TAM DAVIS

Art Unit

1642

-- The MAILING DATE of this communication appears on the cover sheet with the correspondence address --

**Period for Reply**

A SHORTENED STATUTORY PERIOD FOR REPLY IS SET TO EXPIRE 3 MONTH(S) FROM THE MAILING DATE OF THIS COMMUNICATION.

- Extensions of time may be available under the provisions of 37 CFR 1.136(a). In no event, however, may a reply be timely filed after SIX (6) MONTHS from the mailing date of this communication.
- If the period for reply specified above is less than thirty (30) days, a reply within the statutory minimum of thirty (30) days will be considered timely.
- If NO period for reply is specified above, the maximum statutory period will apply and will expire SIX (6) MONTHS from the mailing date of this communication.
- Failure to reply within the set or extended period for reply will, by statute, cause the application to become ABANDONED (35 U.S.C. § 133).
- Any reply received by the Office later than three months after the mailing date of this communication, even if timely filed, may reduce any earned patent term adjustment. See 37 CFR 1.704(b).

**Status**

- 1) ☒ Responsive to communication(s) filed on 25 November 2002.
- 2a) ☐ This action is **FINAL**. 2b) ☒ This action is non-final.
- 3) ☐ Since this application is in condition for allowance except for formal matters, prosecution as to the merits is closed in accordance with the practice under *Ex parte Quayle*, 1935 C.D. 11, 453 O.G. 213.

**Disposition of Claims**

- 4) ☒ Claim(s) 26-63 is/are pending in the application.
- 4a) Of the above claim(s) 29-36, 38-55 and 57-60 is/are withdrawn from consideration.
- 5) ☐ Claim(s) \_\_\_\_\_ is/are allowed.
- 6) ☒ Claim(s) 26-28, 37, 56 and 61-63 is/are rejected.
- 7) ☐ Claim(s) \_\_\_\_\_ is/are objected to.
- 8) ☐ Claim(s) \_\_\_\_\_ are subject to restriction and/or election requirement.

**Application Papers**

- 9) ☐ The specification is objected to by the Examiner.
- 10) ☐ The drawing(s) filed on \_\_\_\_\_ is/are: a) ☐ accepted or b) ☐ objected to by the Examiner.
- Applicant may not request that any objection to the drawing(s) be held in abeyance. See 37 CFR 1.85(a).
- 11) ☐ The proposed drawing correction filed on \_\_\_\_\_ is: a) ☐ approved b) ☐ disapproved by the Examiner.
- If approved, corrected drawings are required in reply to this Office action.
- 12) ☐ The oath or declaration is objected to by the Examiner.

**Priority under 35 U.S.C. §§ 119 and 120**

- 13) ☐ Acknowledgment is made of a claim for foreign priority under 35 U.S.C. § 119(a)-(d) or (f).
- a) ☐ All b) ☐ Some \* c) ☐ None of:
1. ☐ Certified copies of the priority documents have been received.
  2. ☐ Certified copies of the priority documents have been received in Application No. \_\_\_\_\_.
  3. ☐ Copies of the certified copies of the priority documents have been received in this National Stage application from the International Bureau (PCT Rule 17.2(a)).
- \* See the attached detailed Office action for a list of the certified copies not received.
- 14) ☐ Acknowledgment is made of a claim for domestic priority under 35 U.S.C. § 119(e) (to a provisional application).
- a) ☐ The translation of the foreign language provisional application has been received.
- 15) ☐ Acknowledgment is made of a claim for domestic priority under 35 U.S.C. §§ 120 and/or 121.

**Attachment(s)**

- 1) ☐ Notice of References Cited (PTO-892)
- 2) ☐ Notice of Draftsperson's Patent Drawing Review (PTO-948)
- 3) ☐ Information Disclosure Statement(s) (PTO-1449) Paper No(s) \_\_\_\_\_
- 4) ☐ Interview Summary (PTO-413) Paper No(s). \_\_\_\_\_
- 5) ☐ Notice of Informal Patent Application (PTO-152)
- 6) ☒ Other: *See Continuation Sheet*.

Continuation of Attachment(s) 6). Other: courtesy copies of prior search reports in 1997.

Art Unit: 1642

### **DETAILED ACTION**

Effective February 7, 1998, the Group Art Unit location has been changed, and the examiner of the application has been changed. To aid in correlating any papers for this application, all further correspondence regarding this application should be directed to Minh-Tam Davis, Group Art Unit 1642.

A request for continued examination under 37 CFR 1.114, including the fee set forth in 37 CFR 1.17(e), was filed in this application after final rejection. Since this application is eligible for continued examination under 37 CFR 1.114, and the fee set forth in 37 CFR 1.17(e) has been timely paid, the finality of the previous Office action has been withdrawn pursuant to 37 CFR 1.114. Applicant's submission filed on 11/25/02 has been entered.

The text of those sections of Title 35, U.S. Code not included in this action can be found in a prior Office action.

Accordingly, claims 26-28, 37, 56, 61-63, SEQ ID NO:9 are examined in the instant application.

### **REJECTION UNDER 35 USC 112, SECOND PARAGRAPH**

Claims 26-28, 37, 56, 61-63 remain rejected under 35 USC 112, second paragraph pertaining to the use of the language "relative" copy number in claim 26, for reasons already of record in paper No: 28.

Applicant argues that the term "relative copy number" is a term well known in the art, as shown in the paragraph on CGH microarrays obtained from a web site. Applicant

Art Unit: 1642

asserts that the term indicates simply that the measurement, while quantitative need not be an absolute measure of copy number.

The recitation of the paragraph on CGH microarrays is acknowledged.

Applicant's arguments in paper No: 33 have been considered but are found not persuasive for the following reasons:

It is noted that in the paragraph on CGH microarrays, the relative copy number in the test sample is recited as compared to the control sample (3<sup>rd</sup> page, first paragraph). In claim 26 however, it is not clear that the relative copy number of a nucleic acid in a sample is relative to what and/or is compared to what.

Further, there is no definition of the term "relative copy number" in the specification, nor in the CGH microarrays paragraph recited by Applicant.

#### **REJECTION UNDER 35 USC 112, FIRST PARAGRAPH, WRITTEN DESCRIPTION**

Claims 26-28, 56, 61-63 remain rejected under 35 USC 112, first paragraph pertaining to lack of a clear written description, for reasons already of record in paper No: 28.

Applicant argues that Applicant were in possession of a probe which hybridizes to SEQ ID NO:9 under the stringent conditions recited in claim 26. Applicant asserts that the Examiner has offered no objective evidence that such numerous unrelated sequences would be detected in the assay.

Applicant's arguments in paper No: 33 have been considered but are found not persuasive for the following reasons:

The claims as written encompass a method for detecting the presence or absence of neoplastic cells having an increased copy number of nucleic acid sequences at chromosome region 20q13.2, using probes with unknown structure and length, provided said probes share a fragment with SEQ ID NO:9 and are capable of hybridizing to SEQ ID NO:9 via said common fragment under the stringent conditions recited in claim 26.

The specification and the claims lack information of the structure or function of the probes used for the claimed method, and thus do not meet the written description requirement.

**REJECTION UNDER 35 USC 112, FIRST PARAGRAPH, ENABLEMENT, NEW  
REJECTION**

Claims 26-28, 37, 56, 61-63 are rejected under 35 U.S.C. 112, first paragraph, as containing subject matter which was not described in the specification in such a way as to enable one skilled in the art to which it pertains, or with which it is most nearly connected, to make and/or use the invention.

Claims 26-28, 37, 56, 61-63 are drawn to a method of detecting in a sample the presence or absence of neoplastic cells having an increased copy number of nucleic acid sequences at chromosome region 20q13.2, comprising contacting a nucleic acid sample with a probe which hybridizes to SEQ ID NO:9, under the stringent conditions recited in claim 26, and detecting the formation of a hybridization complex to determine the relative copy number of a nucleic acid in the chromosome region 20q13.2, thereby

Art Unit: 1642

identifying the presence or absence of neoplastic cells having an increased copy number of nucleic acid sequences at the chromosomal region 20q13.2.

The specification discloses that using the minimal chromosomal region probe RMC20C001 which is within the chromosome region 20q13.2, an increased level of DNA amplification in the region encompassed by said probe is consistently detected in breast cancers (p.49-50). The specification also discloses that the RMC20C001 probe defines a region of 1.5 Mb within the chromosome region 20q13.2 (p.52, last paragraph). The specification further discloses that SEQ ID NO:9 represent the 2Kb promoter region of zinc finger amplified in breast cancer (ZABC-1), and that this gene maps to the core of the 20q13.2 amplicon and is overexpressed in primary tumors and breast cancer cell lines (p.21, lines 11-15).

No data however is found in the specification concerning the detection of an increased copy number of the gene comprising SEQ ID NO:9. Further, it is not clear whether overexpression of the gene comprising SEQ ID NO:9 is referred to gene amplification or RNA amplification, which are independent from each other .

One cannot extrapolate the teaching of the specification to the scope of the claims because although the 1.5 Mb RMC20C001 probe detect DNA amplification in this region, the RMC20C001 probe spans a very large region of 1.5 Mb which comprises numerous genes that are unrelated to the gene comprising SEQ ID NO:9, which comprises of only a 2Kb sequence, and because it is well known in the art that amplification or regulation of different genes is independent of each other. In other words, it is unpredictable which genes in the 1.5 Mb region are amplified and detected

Art Unit: 1642

by the RMC20C001 probe. Thus one cannot predict that detection of the presence of SEQ ID NO:9 in a sample would detect an increased copy number of the gene comprising SEQ ID NO:9, and it is not clear how the detection of the presence of SEQ ID NO:9 would determine the relative copy number of a nucleic acid in the chromosome region 20q13.2, and thereby indentifying the presence of neoplastic cells.

Further, no specific probes are recited in the claims for use in the detection of SEQ ID NO:8. One would have expected that using any probe, non-related nucleic acid sequences, which share some similarity with SEQ ID NO:9, could be detected, thus could effect the total level of DNA detected. For example, the claimed method would detect 1) a sequence which is 88% similar to SEQ ID NO:9, as taught by Morris et al, 1991 (MPSRCH search report, 1997, us-08-731-499-05.rge, pages 1-2, of record, a courtesy copy of which is enclosed) 2) a sequence which is 84% similar to SEQ ID NO:9, as taught by Ionov Y et al, 1994 (MPSRCH search report, 1997, us-08-731-499-01.rng, page 2, of record, a courtesy copy of which is enclosed) and 3) a sequence which is 77% similar to SEQ ID NO:9, as taught by Beach DH et al, 1993 (MPSRCH search report, 1997, us-08-731-499-01.rng, pages 1-2, of record, a courtesy copy of which is enclosed), wherein these unrelated sequences are not necessarily amplified. In other words, using any probe, it is unpredictable that one could detect an increase in the gene copy of SEQ ID NO:9 in cancer as compared to normal control, due to possible interference by other unrelated sequences that are also detected.

In view of the above, it would have been undue experimentation for one of skill in the art to practice the claimed invention as broadly as claimed.



## **REJECTION UNDER 35 USC 112, FIRST PARAGRAPH, SCOPE**

1. If Applicant could overcome the above 112, first paragraph rejection, claims 26-28, 56, 61-63 are still rejected under 35 USC 112, first paragraph, pertaining to lack of enablement for a method for detecting the presence or absence of “any neoplastic cell” having an increased number of “any nucleic acid sequence” at chromosome region 20q13.2, for reasons already of record in paper No: 28.

Applicant argues that that the Examiner implicitly reads a limitation into the claims that is not present. Applicant asserts that the language “any neoplastic cells” or “any nucleic acid sequences” does not exist in claim 26, and invites the Examiner to identify the language “any neoplastic cells” or “any nucleic acid sequences” in claim 26.

Applicant asserts that genes other than those identified in the present claims may also be amplified at 20q13.2 in neoplastic cells is simply irrelevant to enablement of the claimed invention.

Applicant asserts that the Examiner has provided no objective basis to support an allegation that performing the presently claimed method will fail to identify amplifications at 20q13.2 in a sample containing cells having such amplification.

Applicant's arguments in paper No: 33 have been considered but are found not persuasive for the following reasons:

It is noted that the language “any neoplastic cells” or “any nucleic acid sequences” does not have to be in the claim 26 for the claim 26 to be reasonably interpreted as encompassing a method for detecting the presence or absence of “any

Art Unit: 1642

neoplastic cell" having an increased number of "any nucleic acid sequence" at chromosomal region 20q13.2, by detecting the hybridization of a probe with SEQ ID NO:9, wherein detecting the formation of a hybridization complex would determine the relative copy number of "any nucleic acid" in chromosomal region 20q13.2, thereby indentifying the presence or absence of any neoplastic cells having an increased copy number of "any nucleic acid sequence" at chromosomal region 20q13.2.

One cannot extrapolate from one example of detection of breast cancer, in which SEQ ID NO:9 is overexpressed, to detection of any cancer having increased copy of number of a nucleic acid sequence which is unrelated to SEQ ID NO:9, provided said nucleic acid sequence is within the chromosome region 20q13.2, because using a probe specific for SEQ ID NO:9 one would not expect to detect other sequences that are structurally unrelated to SEQ ID NO:9, but are within the chromosome region 20q13.2. Further, although breast cancer has overexpression of SEQ ID NO:9, it is unpredictable that any other neoplastic cell that has an increased copy number of nucleic acid sequences at chromosome region 20q13.2, wherein said nucleic acid sequences are different than SEQ ID NO:9, would also have an increased copy number of SEQ ID NO:9, because different cancers have different etiology, and mechanisms of carcinogenesis, and because the role of SEQ ID NO:9 in any cancer development is not known.

2. If Applicant could overcome the above 112, first paragraph rejection, claims 26-28, 56, 61-63 are still rejected under 35 USC 112, first paragraph pertaining to lack of enablement for a method for detecting in "any sample", the presence or absence of

Art Unit: 1642

neoplastic cells having an increased number of nucleic acid sequences at chromosome region 20q13.2, for reasons already of record in paper No: 28.

Applicant argues that in a sample lacking neoplastic cells with an amplification at 20q13.2, the assay will be negative, i.e. the assay will report the absence of neoplastic cells in the subject sample as recited in the preamble of the claim.

Applicant's arguments in paper No: 33 have been considered but are found not persuasive for the following reasons:

The claims encompass a method for detecting the presence of neoplastic cells in any sample. However, it is unpredictable that any cancer sample, or any cancer tissue would have an increased copy of SEQ ID NO:9, because different cancers have different etiology, and mechanisms of carcinogenesis, and because the role of SEQ ID NO:9 in any cancer development is not known.

Further, there is no use for the claimed detection of the absence of neoplastic cells in a sample, e.g. in a hair sample.

In view of the above, it would have been undue experimentation to practice the claimed invention as broadly as claimed.

Any inquiry concerning this communication or earlier communications from the examiner should be directed to MINH-TAM DAVIS whose telephone number is 703-305-2008. The examiner can normally be reached on 9:30AM-4:00PM.

If attempts to reach the examiner by telephone are unsuccessful, the examiner's supervisor, ANTHONY CAPUTA can be reached on 703-308-3995. The fax phone


Art Unit: 1642

numbers for the organization where this application or proceeding is assigned are 703-872-9306 for regular communications and 703-872-9307 for After Final communications.

Any inquiry of a general nature or relating to the status of this application or proceeding should be directed to the receptionist whose telephone number is 703-308-0916.

MINH TAM DAVIS

February 4, 2003

  
ANTHONY C. CAPUTA  
SUPERVISORY PATENT EXAMINER  
TECHNOLOGY CENTER 1800



(TM)

Release 2.1D John F. Collins, Biocomputing Research Unit.

Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.

Distribution rights by IntelliGenetics, Inc.

WATERMAN n.a. n.a. database search, using Smith-Waterman algorithm

Run on: Sun Jun 29 03:22:08, 1997 Maspar time 1852.43 Seconds  
1200.504 Million cell updates/sec

Tabular output not generated.

Title: &gt;US-08-731-499-9

Description: (7201-9200) from US08731499.seq (5 of 6)

Perfect Score: 2086

N.A. Sequence: 7201 TATTACCTCATATTCATCT.....AAGCAGACCCCTCTGGGCG 9200

(Comp: TATATGGAGTATAGTAGA.....TTCGTCTGGGAGGACCGG)

Scoring table:

Gap 6

Nmatch: STD Dbase 0; Query 0;

Searched: 33249 seqs, 555961234 bases x 2

Post-processing: Minimum Match 0;

Listing first 45 summaries

Database: emb1-new11

1: BCT1 2: FUN 3: GEN 4: HUM1 5: HUM2 6: HUM3 7: INV1 8: INV2

9: INV3 10: INV4 11: INV5 12: INV6 13: INV7 14: ORG 15: MAM

16: VRT 17: PLN 18: PRO1 19: PRO2 20: ROD 21: SYN 22: UNC

23: VIR1 24: VIR2

Database: genbank97

25: BCT1 26: BCT2 27: BCT3 28: BCT4 29: BCT5 30: BCT6 31: BCT7

32: BCT8 33: BCT9 34: GEN1 35: GEN2 36: HTG 37: INV1 38: INV2

39: INV3 40: INV4 41: INV5 42: INV6 43: INV7 44: INV8 45: INV9

46: MAM1 47: MAM2 48: MAM3 49: VRT1 50: VRT2 51: VRT3 52: PAT1

53: PAT2 54: PAT3 55: PAT4 56: PHG 57: PLN1 58: PLN2 59: PLN3

60: PLN4 61: PLN5 62: PLN6 63: PLN7 64: PLN8 65: PLN9 66: PLN10

67: PRI1 68: PRI2 69: PRI3 70: PRI4 71: PRI5 72: PRI6 73: PRI7

74: PRI8 75: PRI9 76: PRI10 77: PRI11 78: PRI12 79: PRI13

80: PRI14 81: ROD1 82: ROD2 83: ROD3 84: ROD4 85: ROD5 86: ROD6

87: ROD7 88: ROD8 89: STR 90: SYN 91: UNA 92: VRL1 93: VRL2

94: VRL3 95: VRL4 96: VRL5 97: VRL6 98: VRL7 99: VRL8 100: VRL9

Database: genbank-new11

101: BCT 102: GEN 103: INV1 104: INV2 105: MAM 106: VRT

107: PHG 108: PLN 109: PRI1 110: PRI2 111: ROD 112: SYN

113: UNA 114: VRL

Database: u-emb148.97

115: part1 116: part2

Statistics: Mean 12.230; Variance 6.274; scale 1.949

Result: Pred. No. is the number of results predicted by chance to have a

score greater than or equal to the score of the result being printed,

and is derived by analysis of the total score distribution.

SUMMARIES

Result: Query Match Length DB ID Description Pred. No.

No. 1

1

1

1

1

1

1


1

1

1

1





\*\*\*\*\*  
Release 2.1D John F. Collins, Biocomputing Research Unit.  
Copyright (c) 1993, 1994, 1995 University of Edinburgh, U.K.  
Distribution rights by Intelligenetics, Inc.  
\*\*\*\*\*  
MPSrch\_nn n.a. - n.a. database search, using Smith-Waterman algorithm  
Run on: Sat Jun 28 22:40:56 1997; MasPar time 213.20 Seconds  
Tabular output not generated. 867.829 Million cell updates/sec  
\*\*\*\*\*  
Title: >US-08-731-499-9  
Description: (1-2000) from US08731499.seq (1 of 6)  
Perfect Score: 2000  
N.A. Sequence: 1 CCATCATATTCTTATTTT.....AAATAGTTTACTAAAGTGT 2000  
Comp: GGTACTATAAGATAAAAA.....TTTATCAATGATTTCACA  
\*\*\*\*\*

Scoring table: TABLE default  
Gap 6  
Nmatch STD : Dbase 0; Query 0  
Searched: 121476 seqs, 46255616 bases x 2  
Post-processing: Minimum Match 0  
Listing first 45 summaries  
Database: n-geneseq26  
1:part1 2:part2 3:part3 4:part4 5:part5 6:part6 7:part7  
8:part8 9:part9 10:part10 11:part11 12:part12 13:part13  
14:part14 15:part15 16:part16 17:part17 18:part18  
19:part19 20:part20 21:part21 22:part22 23:part23  
Statistics: Mean 9.913; Variance 8.353; scale 1.187  
Pred. No. is the number of results predicted by chance to have a  
score greater than or equal to the score of the result being printed,  
and is derived by analysis of the total score distribution.

SUMMARIES					
Result No.	Score	Query Match	Length	ID	Description
C 1	135	6.8	3158	9	Q53212 Human cyclin D3 promo
C 2	135	6.8	3158	5	Q31280 Cyclin D3 promoter.
C 3	121	6.1	283	11	Q63862 AP2 sequence obtd. by
C 4	120	6.0	7849	16	Q94109 hML genomic DNA.
C 5	118	5.9	1618	7	Q46558 Human cytokine synthe
C 6	118	5.9	1618	2	Q10207 p15C insert containi
C 7	118	5.9	6511	14	Q95493 Human Cdn-2 DNA.
C 8	117	5.8	321	8	Q59208 Human brain Expressed
C 9	117	5.8	321	8	Q39796 Expressed Sequence Ta
C 10	116	5.8	335	8	Q60863 Human brain Expressed
C 11	117	5.8	2600	1	N90029 Human interleukin-1 r
C 12	117	5.8	2600	12	Q37364 Human IL-1 receptor g
C 13	117	5.8	2600	1	N90118 cDNA of human interle
C 14	117	5.8	2963	8	Q49933 IL-1R cDNA.
C 15	117	5.8	3234	15	Q92781 Human thymopoietin ge
C 16	115	5.8	11531	9	Q54222 BSSL/CEL Gene.
C 17	117	5.8	17327	7	Q44278 Serglycin - proteogly

18 114 5.7 4823 22 T37384 Human thrombopoietin 1.79e-38  
19 114 5.7 4823 17 T03943 Human thrombopoietin 1.79e-38  
20 114 5.7 4823 16 T04051 Sequence encoding hae 1.79e-38  
21 112 5.6 2435 23 T33155 Tissue plasminogen ac 1.59e-37  
22 112 5.6 2888 1 Q03743 Human SKI related gen 1.59e-37  
23 112 5.6 6905 15 Q92779 Human thymopoietin co 1.59e-37  
24 111 5.6 8363 15 Q92408 Human cyclin A gene. 4.75e-37  
25 112 5.6 17327 7 Q44278 Serglycin - proteogly 1.59e-37  
26 112 5.6 22481 23 T11658 PEDF full length sequ 1.59e-37  
27 110 5.5 2649 6 Q35034 DNA fragment contgs. A 1.41e-36  
28 110 5.5 7620 6 Q3286 Glucocerebrosidase ge 1.41e-36  
29 110 5.5 10884 23 T33758 Control region isolat 1.41e-36  
30 109 5.4 308 8 Q60826 Human brain Expressed 4.20e-36  
31 109 5.4 743 2 N70812 Sequence encoding hum 4.20e-36  
32 109 5.4 2660 3 N30032 Sequence of gene for 4.20e-36  
33 108 5.4 4382 2 Q12759 P40 genomic DNA. 1.25e-35  
34 108 5.4 6063 6 Q37205 Delta-amino levulinat 1.25e-35  
35 108 5.4 10897 19 T09187 Mutu putative oncogen 1.25e-35  
36 109 5.4 13585 17 T11549 Tumour rejection anti 4.20e-36  
37 109 5.4 30967 23 T32454 Calpain large subunit 4.20e-36  
38 106 5.3 366 8 Q60353 Human brain Expressed 1.10e-34  
39 107 5.3 1777 12 Q79355 Sequence of the exten 3.70e-35  
40 107 5.3 2320 4 Q28657 glut4 promoter/enhanc 3.70e-35  
41 107 5.3 2339 2 Q10956 Encodes human 75kD TN 3.70e-35  
42 107 5.3 2425 18 T11027 DNA encoding the huma 3.70e-35  
43 107 5.3 2425 20 T10283 Gene for RNA componen 3.70e-35  
44 106 5.3 5359 17 T12251 Cytochrome P450 isoen 1.10e-34  
45 107 5.3 9272 12 Q79353 Human genomic clone h 3.70e-35

ALIGNMENTS

RESULT 1  
ID Q53212 standard; DNA; 3158 BP.  
AC Q53212;  
DT 22-JUN-1994 (first entry)  
DE Human cyclin D3 promoter.  
KW D-type; mammalian; CLN protein; protein deficiency; cell cycle start;  
OS Yeast; complement; ds.  
KW Homo sapiens.  
FH Key Location/Qualifiers  
FT misc\_feature 3158..3158  
FT /\*tag= a  
FT /note= "Initiation ATG codon"  
PN WO9324514-A  
PD 09-DEC-1993  
PF 25-MAY-1993; U05000.  
PR 26-MAY-1992; US-888178.  
PI (MITO-) MITOTIX.  
PI Beach DH.  
DR WPI; 93-405720/50.  
PT New D-type mammalian cyclin - replaces CLN-type protein needed  
PT for cell start in budding yeast and is detected by antibodies or  
PT hybridisation in biological samples to determine abnormal cell  
PT division  
PS Disclosure: Fig 13; 108pp; English.  
CC The sequence is that of human cyclin D3 promoter.  
SQ Sequence 3158 BP; 952 A; 674 C; 722 G; 810 T;  
Query Match 6.8%; Score 135; DB 9; Length 3158;  
Best Local Similarity 77.0%; Pred. No. 1.64e-48;  
Matches 221; Conservative 0; Mismatches 62; Indels 4; Gaps 2;  
Db 1653 gpcgggagcgggtgctcagcgtatccagcagcactttggagcgagaccgagcga 1712  
Cp 293 GGACGGGTATAGTGGCTCACACCTATATCTCCCATGCTTTGGGAGCCGGAAGTGGTGA 234  
Db 1713 tcac--gaggttcagggttcagactagctggccacatagtgaaacccctctctacg 1770  
Cp 233 TCACATAGGCCAGGAGTGGAGACCGCCGCGCCAAACATGGTGAACCCCTTATCTGCT 174  
Db 1771 aaaaatacaaaaattagtcaggcatggtggtgctgtagtcctccagctactccgggaa 1830  
|||||

8/13/97

✓	173	AAATAACAAAAATTAGCTGGCGATGTAATACACACTGTAATCCAGTATTTGGAA	114
Db	1831	ttgcttgaacccggaggttgaggttgcagtagcgcacatcgccacactgcactccagc	1890
Cp	113	TCACTTGAACCCAGAGGTGGAGTGTGCAGTGAGCCAAAGATCGCACACCTGG--TCAGC	56
Db	1891	ttgagcacacagtagtgacttcgtctcaaaaaaataaaaaaataaaaaa	1937
Cp	55	CTGGGCAACAGACGAAGTCTCCCTCTCCGCCCAAAAAAATAAGAA	9

2  
RESULT  
ID Q31880 standard; DNA; 3158 BP.  
AC Q31880;  
DE 22-APR-1993 (first entry)  
DE Cyclin D3 promoter;  
KW Cyclin; D1; D2; D3; promoter; human; liver; genomic library; clone;  
KW upstream; exon; intron; neural; pCYCD1-H12; mutant; yeast; strain;  
KW CLN; cyclin; gene; CLN 1; CLN 2; human; glioblastoma; cDNA library;  
KW expression vector; PADNS; transformant; pCYCD1-21; pCYCD1-19; HeLa;  
KW as.  
PS Homo sapiens.  
PN WQ9220756-A.  
PN 26-NOV-1992.  
PF 18-MAY-1992; U04146.  
PF 16-MAY-1991; US-701514.  
PA (COLD-) COLD SPRING HARBOR LAB.  
PI Beach DH;  
PI WPI; 92-415774/50.  
DR Recombinant mammalian D-type cyclin - replaces a CLN-type protein  
PT essential for cell start in budding yeast, its antibodies and  
PT probes being useful in detecting D-type cyclin in biological  
PT samples  
PS Disclosure; Fig 13; 75pp; English.  
CC The sequences given in Q31878-80 represents the cyclin D1 to D3  
CC promoters. These sequences were identified during the isolation of  
CC the D-type cyclin cDNAs from a normal human liver genomic library.  
CC A mutant yeast strain in which two of the three CLN cyclin genes  
CC (CLN 1 and CLN 2) were inactivate and expression of the third was  
CC conditional, was used to identify human cDNA clones that rescue yeast  
CC from CLN deficiency. A human glioblastoma cDNA library carried in a  
CC yeast expression vector (PADNS) was introduced into a mutant yeast  
CC strain. Two yeast transformants (pCYCD1-21 and pCYCD1-19) which grew  
CC despite the lack of function of all three CLN genes and were not  
CC revertants, were identified and recovered in E. coli. These two  
CC clones were shown to be independent clone representing the same gene.  
CC A HeLa cDNA library was screened for a full length cDNA clone using the  
CC 1.2 kb insert of pCYCD1-21 as a probe. The sequence isolated by this  
CC method was pCYCD1-H12 (see also Q31873). Degenerate probes and  
CC primers were designed using the D1 gene sequence. These primers  
CC and probes were used in the isolation of the cyclin D2 and D3 genes.  
CC See also Q31874-5. The cyclin D1 cDNA clone was used to screen a  
CC liver genomic library resulting in the identification of three  
CC positive clones. These clones were shown to correspond to the  
CC upstream promoter region and a 198 bp exon, followed by an intron of  
CC cyclin D1. Human cyclin promoters D2 and D3 were isolated in the same  
CC manner. Cyclin D1 has been shown to be expressed differentially in  
CC different cell types, with expression being highest in cells of neural  
CC origin.  
SQ Sequence 3158 BP. 952 A; 674 C; 722 G; 810 T;

Query Match	6.8%	Score 135;	DB 5;	Length 3158;
Best Local Similarity	77.0%;	Pred. No. 1.64e-48;		
Matches	221;	Conservative	0;	Mismatches 62; Indels 4; Gaps 2;
Db	1653	gcccgggaacggtgctcacgcgtgtaatcccgacacatttggagcgccgagaccgcgga	1712	
Cp	293	GGACGGGTATAGTGGCTCACCTATACPCCCAATGCTTTGGGAAGCCGAAGTGGGTGA	234	
Db	1713	tcac--gaggtcaggggttcagactagcctggccacaatagtgaaaccccatctctacg	1770	
Cp	233	TCACATGAGGCCAGGAGTTGGAGACAGCGCTGGCCAAACATGTTGAACCCCTTATCTGCT	174	

[illegible]

RESULT 3

ID Q63862 standard; cDNA; 283 BP.  
AC Q63862;  
DE 29-JAN-1995 (first entry)  
DT AP2 sequence obtd. by PCR for tumour specific DNA.  
DE Arbitrary Primers; AP-PCR; amplification; tumour cells; cancer;  
KW Arbitrary Primers; AP-PCR; amplification; tumour cells; cancer;  
KW Insertions; deletions; ss.  
OS Synthetic.  
PN W09411531-A.  
PD 26-MAY-1994.  
PP 12-NOV-1993; U10904.  
PR 13-NOV-1992; US-975737.  
PA (CALB-) CALIFORNIA INST BIOLOGICAL RES.  
PI Ionov Y, Malkhosyan S, McClelland M, Peinado MA;  
PI Perucho M, Welsh J;  
DR WPI: 94-183529/22.  
PT Identification of tumour cells - by analysing DNA to determine  
PT whether insertions or deletions have occurred in reiterated  
PT sequences  
PS Disclosure; Page 52; 67pp; English.

CC	The sequence was obtd.	by PCR with arbitrary PCR primers used to
CC	detect insertions or deletions	in DNA sequences. Such mutations are
CC	markers of cancer so such primers can be used	in the diagnosis of
CC	cancer, esp. colorectal, stomach or pancreatic tumours.	
CC	See also Q63837-63.	
SQ	Sequence 283 BP:	63 A; 77 C; 94 G; 49 T;
	Query Match	G-114 Score 121; DB 11; Length 283;
	Best Local Similarity	84.58; Pred. No. 8,37e-42;
	Matches 148; Conservative	0; Mismatches 21; Indels 0; Gaps 0;
Ddb	5	ggcggtggtgcacacctgtaatccacagactttggaggcggagtgggtggatacac 64
Cp	289	GGGTATAGTGGCTCACACCTATACTCCCAATGCTTTGGGAACCGAATGGGTGGATCACC 230
Ddb	65	ctgaaggtcaggaggttcagaccagcctgcccaacatggtgaaaccccgcctctactaaaa 124
Ccp	229	ATGAGGCCAGGAGTGAGACACCGAGCTGCACAACATGGTGAACCCCTTATCTGCTAAAA 170
Ddb	125	atacaaaaattagccggcgctggttggcgcgcgctgttaatccccagctactcggga 179
Ccp	169	ATACAAAATTAGCTGGGCATGGTAAATACACACCTGTATCCACGATTTTGGGA 115

RESULT	4	
ID	Q94109 standard; DNA; 7849 bp.	
AC	Q94109;	
DT	22-FEB-1996 (first entry)	
DE	hML genomic DNA.	
KW	Human; thrombopoietin; TPO; mpl ligand; hML; fragment polypeptide; megakaryocytopoietic cytokine receptor; thrombopoietic signal; EPO-domain fragment; erythropoietin; hEPO; haematopoietic cell; megakaryocyte; thrombocytopenia; myeloproliferative disease; inflammatory thrombocytosis; iron deficiency; EPO; platelet; red blood cell; progenitor; hML-2; ss.	
OS	Homo sapiens.	
PH	Key	Location/Qualifiers
Tt	prim_transcript	ll66..7289
Tt	/tag= a	
Tt	exon	ll61..1232